

Revista Brasileira de Zootecnia



All the contents of this journal, except where otherwise noted, is licensed under a Creative Commons Attribution License. Fonte:

https://www.scielo.br/scielo.php?script=sci_arttext&pid=S1516-35982013001000007&lng=en&tln g=en. Acesso em: 05 ago. 2020.

REFERÊNCIA

CARDOSO, Maximiliano Tadeu Memória et al. Performance and carcass quality in three genetic groups of sheep in Brazil. **Revista Brasileira de Zootecnia**, Viçosa, v. 42, n. 10, p. 734-742, out. 2013. DOI: <https://doi.org/10.1590/S1516-35982013001000007>. Disponível em: https://www.scielo.br/scielo.php?script=sci_arttext&pid=S1516-35982013001000007&lng=en&tln g=en. Acesso em: 05 ago. 2020.



Performance and carcass quality in three genetic groups of sheep in Brazil

Maximiliano Tadeu Memória Cardoso¹, Aline Vieira Landim², Helder Louvandini³,
Concepta McManus^{1,4}

¹ Faculdade de Agronomia e Medicina Veterinária, Universidade de Brasília, Brasília, DF, Brasil.

² Departamento de Zootecnia, Universidade Estadual Vale do Acaraú, Sobral, CE, Brasil.

³ Centro de Energia Nuclear na Agricultura, Piracicaba, SP, Brasil.

⁴ Departamento de Zootecnia, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brasil.

ABSTRACT - Three different genetic groups were evaluated for growth up to slaughter and carcass quality for kill out, commercial cuts, non-carcass components, as well as carcass morphometric measurements. Pure Santa Ines (SI), ½ Ile de France × ½ Santa Ines (I × SI) and ½ Texel × ½ Santa Ines (T × SI) lambs were weighed weekly until they reached predetermined slaughter weight (30, 35, 40 and 45 kg). In both creep feeding and fattening the animals from crosses were superior to the Santa Ines hair sheep in terms of daily weight gain and weaned in less time. Crossbred animals also showed better performance than purebred Santa Ines in terms of commercial cuts and carcass weights. Correlations showed that fat depth and eye muscle area decreased with an increase in morphometric measures, indicating that larger animals are possibly later developing in terms of carcass finishing compared with smaller animals. The optimal slaughter weight is approximately 35 kg.

Key Words: creep feeding, cross-breeding, fattening, lambs

Introduction

Commercial sheep production for meat in Brazil is still in the early stages of development, with live weight determining slaughter age. According to Juárez et al. (2009), consumer markets demand minimum weights for carcass cuts thereby avoiding slaughter of animals not in a finished condition. The determination of ideal slaughter weight varies depending on regions and countries and affects carcass quality traits such as organoleptic quality, fat deposition as well as subjective traits (Landim et al., 2011a,b).

Live weights and subjective assessments of condition or conformation are commonly used by farmers to select lambs with the best potential carcass quality characteristics (Lambe et al., 2008). Live weight at slaughter may be altered by the way the animal is slaughtered including the effects of fasting time and conditions, as well as time of year and transport (Brasal & Boccard, 1977; Pérez et al., 2007).

Quantitative carcass measurements such as relative and actual weights are important as these are criteria used to evaluate animal productivity (Zundt et al., 2001). The finished carcass is the result of a biological process affected by genetic, environmental and management factors, and carcasses differ in quantitative and qualitative factors (Osório & Osório, 2001).

Sheep breeding in Brazil is based mainly on the Santa Ines, which is a hair breed with poor carcass quality and slow growth rate. The use of terminal sire meat breeds with

higher growth rates and feeding strategies are resources which have been used to reduce slaughter age and improve carcass quality (Macedo et al., 2000). Crossbreeding can also lead to animals with higher growth rates, better conformation and better meat quality in lambs (Cardellino, 1989).

Slaughter weight for hair sheep in center-west Brazil is generally fixed at 30 kg (Paim et al., 2011), which is significantly lower than that used for terminal sire breeds. These same authors found that the most profitable live weight for slaughter was 45 kg. New studies on the definition of slaughter age and reduction of costs are necessary. This study aims to evaluate the performance of Santa Ines lambs as well as crosses with Texel and Ile de France, slaughtered at different weights, in creep feeding and fattening in tropical conditions.

Material and Methods

This experiment was carried out in the Sheep management Center of the Água Limpa farm of Universidade de Brasília, in the Federal District, Center West region of Brazil. The climate is classified as Aw by Köppen, with two distinct seasons (rainy and dry) and a mean annual temperature of 23 °C. Mean annual precipitation is 1330 mm and relative humidity 66%.

Entire male and female lambs from three genetic groups - pure Santa Inês (SI) and crossed products (I × SI and T × SI) - were slaughtered at four different live weights (30, 35, 40 and 45 kg), with six animals in each group.

After birth, lambs were individually identified, creep-fed and weaned at 18 kg live weight. After weaning they were dewormed and housed until reaching their randomly predefined slaughter weight.

The animals received coast cross (*Cynodon dactylon*) hay (dry matter (DM): 873 g/kg; crude protein: 68 g/kg; neutral detergent fiber (NDF): 783 g/kg; acid detergent fiber (ADF): 337 g/kg; ether extract (EE): 15 g/kg; ash (A): 14 g/kg) and a concentrate (DM: 909 g/kg; crude protein: 188 g/kg; NDF: 325 g/kg; ADF: 337 g/kg; EE: 32 g/kg; A: 74 g/kg) of corn (561.4 g/kg), soybean meal (295.7 g/kg), wheat flour (102.9 g/kg) and a mineral vitamin nucleus (4 g/kg), as well as a mineral mixture (Ovinofós®) and water *ad libitum*. The diet was offered twice daily at 4 g/kg live weight, with 70 kg/100 kg concentrate and 30 kg/100 kg hay.

Animals were weighed weekly and slaughtered at their predetermined weight. Body measurements were carried out on the day prior to slaughter and obtained according to Santana (2001) and Osório et al. (1998). These included:

Shoulder height (SH): measured using a hipometer, as the vertical distance between the highest point of the shoulder and the soil with the animal standing square; Body length (BL): distance between base of the tail and base of the neck; Heart girth (HG): circumference of the trunk posterior to the shoulder blades; Cannon bone perimeter: measured in the medial portion of the metacarpal bone; Body score (BS): measured by palpation to measure the quantity of fat between second and fifth lumbar vertebrae (L2 to L5), in the region of the sternum and at the base of the tail. The scores are subjective, varying from 1 to 5, where 1 = very thin and 5 = very fat.

Animals were weighed before fasting to obtain live weight (LW). The animals were weighed before slaughter after fasting without water or food for 16 hours to obtain slaughter weight (SW). Animals received an electric shock to desensitize them, then the jugular vein and carotid arteries were cut and animals bled out. The animals were skinned, head was removed and the skin was weighed. The abdomen and thorax were cut along the median line and viscera were removed and weighed. Thoracic (lung, heart and trachea) organs were weighed separately from those of the abdominal cavity (liver and kidneys). Digestive tract contents were removed and weighed and empty body weight was calculated (Silva Sobrinho, 2001).

The carcasses were weighed to obtain hot carcass weight (HCW), and hot carcass killout was calculated ($HCK = HCW/SW \times 100$). The carcasses were then refrigerated at 4 °C for 24 hours and weighed again to give cold carcass weight (CCW), loss during refrigeration

($LR = HCW - CCW/HCW \times 100$) and commercial or cold carcass killout ($CK = CCW/SW \times 100$).

Carcass traits were evaluated using the system proposed by Osório et al. (1998). Fat cover (FC) was evaluated subjectively by the quantity and distribution on the outside of the carcass varying from 1 (very thin) to 5 (very fat). Carcass length (CL) was measured as the distance between the base of the tail and base of the neck.

The carcass was divided in half along the back bone and the left half was weighed (CWH) along with six commercial cuts: leg, shoulder, back, rib/belly and neck. True killout (TK) of hot carcass was calculated using the method proposed by Osório et al. (1998).

The experimental design was completely randomized in a factorial arrangement with 3 genetic groups and 4 slaughter weights. Data were collected using procedures MIXED (General Linear Model), CORR (Correlation) and REG (Regression) of live weight on carcass traits on Statistical Analysis System (SAS, version 9.1). Means were tested using Duncan test as $\alpha = 0.05$. Only two significant interactions were found for slaughter weight * sex for head and intestine weights. No other significant interactions were found between factors and so are not discussed here.

Results and Discussion

Genetic group did not affect birth weight or weight at weaning, fixed at 18 kg (Table 1). Birth weight was 3.04, 2.94 and 2.95 kg for Texel × Santa Ines, Ile de France × Santa Ines and pure Santa Ines animals, respectively.

Texel × Santa Ines and Ile de France × Santa Ines showed higher weight gain up to slaughter, while the Santa Ines had the lowest (Table 2). No significant differences were found between the growth rates of lambs in the different groups slaughtered at different weights.

According to Moreno et al. (2010), specialized breeds for meat production such as Ile de France, Texel, Suffolk, Dorper and Hampshire have high weight gains, which lead to early weaning and earlier slaughter. Crossbreeding with exotic breeds such as Texel and Ile de France is increasing and rearing climates are vastly different from those found in regions from where many of the breeds used today in production systems were developed. These include type of forage available (C3 vs C4), temperatures, humidity, solar radiation and types of feed available.

Genetic group affected animal production, as Ile de France and Texel are terminal meat breeds, selected for increased growth rate while the Santa Ines is a naturalized Brazilian hair sheep. These crosses benefit from hybrid vigor. The Texel × Santa Ines showed a 31% higher daily

gain than Santa Ines, while Ile de France \times Santa Ines grew 2% faster.

Males showed higher weight gain than females, averaging 143 and 122 g/animal/day, respectively over all slaughter weight classes. Males therefore grew 19% faster than females. This was reflected in days from weaning to slaughter (DWS); males were slaughtered earlier than females (26 days or 21% less time). No significant differences were found in growth rates or days to slaughter between lambs born as single or twins, although there was a tendency for single-born lambs to grow faster. Studies show that differences between genetic groups, sexual condition, diet, among others, may influence lamb performance (Arnold & Meyer, 1988; Ribeiro et al., 2002).

According to Pinheiro et al. (2007), most sheep meat commercialized in large urban centers in Brazil does not contain information such as breed, origin, feed information, sex, or age of animal. The consumer therefore may acquire meat of diverse qualities with different proportions of muscle, bone and fat, which may limit consumption and

commercialization of this type of animal. There is a need therefore to describe meat from different sources, breeds, sexes and slaughter weights reared in differing Brazilian ecosystems so the consumer can better chose meat of their preference.

Overall, the means found were 61.86, 79.23 and 64.96 cm for BL, HG and SH, respectively (Table 3). Slaughter weight was significant for all morphological traits and generally showed a linear effect except for cannon bone perimeter, where it was quadratic. Thoracic perimeter increased with slaughter weight, but no differences between sexes were seen for this trait. Females were longer, fatter, with finer bones. Animals from the genetic group T \times SI were longer and had a higher girth than the other groups, while the crossbred animals had larger bones than the SI animals.

The means for morphometric measures in pure Santa Inês were similar to those found by McManus & Miranda (1997), who, studying Santa Ines, found HG and BL to be 82.34 and 64.94 cm, respectively. Landim et al. (2007) found for BL, HG and SH 61.90, 72.71 and 63.00 cm, respectively, for lambs raised at pasture and slaughtered at 30 kg. Similar results were found in the present study with 30 kg lambs with mean values of BL 57.64, HG 74.18 and SH 60 cm. Mendonça et al. (2003) found a mean thoracic circumference of 77.84 cm for Corriedale and Ideal lambs, while Santana et al. (2001) found that thoracic circumference was highly correlated with body weight in young Santa Ines lambs.

Rosa et al. (2002) found no significant differences between biometric measures between crossbred Ideal \times Ile de France or Texel lambs, in different feeding systems but slaughtered at the same weight. It was supposed that biometric measures are not influenced by feeding system but genetic group and slaughter weight, as seen here.

Table 1 - Least squared means for pure and crossbred lambs in creep feeding

Source	Birth weight (kg)	Daily growth rate (g/day)	Days to weaning (days)
Genetic group			
T \times SI	3.04	222.57a	67.62a
I \times SI	2.94	204.66a	76.45a
SI	2.95	191.60b	81.08b
Sex			
Male	3.12	201.8	76.75
Female	2.79	201.2	77.69
Lambing type			
Single	3.02	207.53a	74.20a
Twin	2.69	175.09b	90.40b

Different letters in the same column are significantly different ($P < 0.05$) using Duncan test.

T \times SI - $\frac{1}{2}$ Texel \times $\frac{1}{2}$ Santa Ines; I \times SI - $\frac{1}{2}$ Ile de France \times $\frac{1}{2}$ Santa Ines; SI - pure Santa Ines.

Table 2 - Days to slaughter and growth rates in fattening by genetic group for lambs slaughtered at different weights

Weight	Days to slaughter				Growth rate in fattening (kg/day)			
	30	35	40	45	30	35	40	45
Genetic group								
T \times SI	-	104a	-	138a	-	0.142c	-	0.182a
I \times SI	76.6a	128b	159a	150b	0.143a	0.113b	0.129a	0.165b
SI	117.8b	138c	164b	177c	0.081b	0.102a	0.113b	0.124c
Sex								
Male	90.2a	123.6a	149.5a	145.6a	0.117a	0.116a	0.135a	0.124a
Female	109.8b	131.7b	174.6b	180.2b	0.101b	0.105b	0.106b	0.109b
Birth type								
Twin	105.0	131	164	189	0.086	0.105	0.115	0.107
Single	98.5	120	156	168	0.112	0.114	0.136	0.115

Different letters in the same column are significantly different ($P < 0.05$) using Duncan test.

T \times SI - $\frac{1}{2}$ Texel \times $\frac{1}{2}$ Santa Ines; I \times SI - $\frac{1}{2}$ Ile de France \times $\frac{1}{2}$ Santa Ines; SI - pure Santa Ines.

Significant differences were found between genetic groups for: HCW, CCW and CWH. The crossbred animals had a higher HCW than the Santa Ines animals, but did not differ among themselves, while the Texel cross showed heaviest CCW and CWH (Table 4).

The results for HCW and CCW are in agreement with Martins (1997) and Landim et al. (2007), who found that crossbred animals showed higher growth potential than purebred Santa Ines when reared at pasture.

With an increase in slaughter weight, carcass weights were higher, as did killout, probably due to a decrease in non-carcass components. Animals slaughtered at 35 and 45 kg did not differ for hot carcass and cold carcass killout, although animals slaughtered at 45 kg had higher fat deposition. In general, the 35 kg slaughter weight showed the best carcass quality.

No significant differences were found between genetic groups for body score. Shoulder height did not vary between

genetic group or sex (Table 3). Animals slaughtered at 30 kg tended to be shorter, thinner and have smaller bones than the other groups. According to Pires et al. (2006), lambs finished indoors and slaughtered at 30 kg live weight had mean hot and cold carcass weights of 15.19 and 14.47 kg, respectively. Higher hot carcass, cold carcass and half carcass weights were found for the crossbred animals, indicating that the Santa Ines has a high potential for use in crossbreeding using specialized meat breeds such as Texel and Ile de France.

No significant differences were found between genetic groups for loss in refrigeration (LR), commercial killout and hot carcass kill out percentages ($P>0.05$). Nevertheless, there was a tendency for Santa Ines sheep to show higher LR and lower carcass killout than other groups.

The values for loss in refrigeration are higher than those found by Osório et al. (2002), who observed hot and cold carcass killout of 46.3 and 42.9 kg/100 kg, respectively, for

Table 3 - Means for morphometric measures (cm) in pure and crossbred lambs

Source of variation	Body length	Body score	Heart girth	Shoulder height	Cannon bone perimeter
Genetic group					
Texel × Santa Ines	65.00a	3.52	83.00a	66.00	9.00a
Ile de France × Santa Ines	62.25b	3.50	80.12b	65.29	8.86a
Santa Ines	60.24b	3.25	77.20c	64.33	8.56b
Sex					
Male	60.89a	3.26a	78.96	65.68	9.07a
Female	62.88b	3.54b	79.52	64.21	8.40b
Slaughter weight (kg)					
30	57.64a	2.82a	74.18a	60.00a	8.31a
35	60.53b	3.20b	76.57b	65.67b	8.30a
40	62.54b	3.54c	79.91c	65.27b	8.91b
45	65.56c	3.87c	84.75d	67.50b	9.34b
Regression	47.51+0.44*LW	0.243+0.088*LW	47.38+0.89*LW	51.83+36*LW	6.6943+0.0016*LW ²

Means with different letters in the same column are significantly different ($P<0.05$), using Duncan test.
LW - live weight.

Table 4 - Mean comparison for carcass traits in purebred and crossbred sheep

Source of variation	Hot carcass weight (kg)	Cold carcass weight (kg)	Half carcass weight (kg)	Loss of weight (kg/100 kg)	Hot carcass killout (kg/100 kg)	Carcass killout (kg/100 kg)	Carcass fat score (1 to 5)
Genetic group							
Tx SI	19.20a	18.68a	9.38a	2.73	46.79	55.42	3.62
I × SI	18.07a	17.22b	8.61b	2.40	46.03	55.42	3.71
SI	17.62b	17.10b	8.58b	4.36	47.10	57.24	3.42
Sex							
Male	17.66	17.17	8.63	2.82	45.86	55.34	3.52
Female	18.47	17.61	8.80	3.86	48.24	57.19	3.61
Slaughter weight (kg)							
30	13.60a	13.20a	6.61a	2.91a	44.47	52.72a	3.27a
35	17.12b	15.83b	7.93b	5.69a	48.08	58.46b	3.47a
40	19.14c	18.66c	9.34c	2.41a	47.82	57.54b	3.36a
45	21.28d	20.84d	10.45d	2.04a	47.25	55.70b	4.00b
Regression	0.19+0.51*LW	-0.11+0.51*LW	0.24+0.45*LW	NS	NS	8.545+0.003*LW	0.243+0.088*LW

Means with different letters in the same column are significantly different ($P<0.05$), using Duncan test.

T × SI - ½ Texel × ½ Santa Ines; I × SI - ½ Ile de France × ½ Santa Ines; SI - pure Santa Ines; LW - live weight; NS - not significant.

crossbred lambs at pasture slaughtered at 33 kg. According to Pinheiro (2006), the fat cover in the carcass, the maturity of the animal, the atmospheric conditions in the meat freezer and time of storage are factors that interfere in the percentage of weight lost during chilling.

Roque et al. (1999) found that the faster the growth rate of the animal, the faster fat is deposited. This was not observed in the present study. Furusho-Garcia et al. (2004) found LR of 4.3 kg/100 kg in Santa Ines lambs slaughtered at 35 kg and 156 days of age, while Osório et al. (2002) found 7.2 kg/100 kg in crossbred Border Leicester × Corriedale and Border Leicester × Ideal lambs.

Studies show differing effects of sex on carcass components. Silva et al. (2000) found heavier shoulder in males than females. Pires et al. (2011) found that sex did not influence the weight of shoulder in males (18.05 kg/100 kg) and females (18.62 kg/100 kg) in Ile de France × Texel lambs. This is due to the 100% direct heterosis obtained in cross products when compared with pure SI animals.

The T × SI cross had the heaviest leg, back and rib of the three genetic groups studied (Table 5). Results show a 15% higher leg weight in the Texel and 6.3% higher in the Ile de France compared with Santa Ines, crossbred animals having more muscle due to the selection undergone for meat production in these breeds. In general, the 35 kg slaughter weight showed the best carcass quality. For the Santa Ines, the best weight was 40 kg, with higher proportions of expensive cuts.

According to Pilar (2002), the proportions of the carcass cuts are an important index for the commercial evaluation of the carcass and have different economic values. Factors such as genetics, diet, slaughter weight, sex among others, are responsible for differences in cuts between carcasses.

Most carcass cuts showed a linear regression on weight at slaughter except for leg measurements, which was quadratic. This may reflect the late maturing of this limb.

The Santa Ines had the lightest skin, head and intestines, the last two not differing from Ile cross animals (Table 6). The differences in skin weight may be due to the fact that SI is a hair breed and the others are wool breeds. Females had heavier abdominal organs and lighter rumen, head and skin than males. Thoracic and abdominal organs are early or intermediate maturing, and so the 30 kg slaughter weight differed from the others.

Heavier lambs here tended to have heavier non-carcass components. Edible viscera can account for up to 5% of the receipts obtained from commercialization of the carcass (Costa et al., 2003). Huidobro & Villapadierna (1992) classified certain parts of the body such as heart, head and kidneys as early maturing and others such as lungs, spleen, small intestine and blood as intermediate, while skin, pancreas, stomach and large intestine are late maturing.

Texel cross lambs had relatively larger muscles (Table 7). Females also had higher fat deposits. Fat depth and rib eye area decrease with an increase in morphometric measures, indicating that larger sized animals were larger developing in terms of carcass finishing than smaller animals.

Rosa et al. (2002) showed that the digestive tract of lambs is late developing and when more time is allowed to slaughter, the reticulum-rumen is more developed. Consequently, its relative contribution to the weight of the animal increases and killout decreases. The Texel here indicates that the rumen weight reflects earlier maturing. The size of the *longissimus dorsi*, which is late maturing, is directly related to body development and quantity of muscle in the carcass (Taylor, 1985).

Table 5 - Mean comparison for commercial carcass cuts in purebred and crossbred lambs in the Federal District, Brazil

Source of variation	Leg (kg)	Leg perimeter (cm)	Back (kg)	Shoulder (kg)	Rib (kg)	Neck (kg)	Belly (kg)
Genetic group							
T × SI	2.91a	37.66a	0.72a	1.51	2.59a	0.91	0.56
I × SI	2.69b	37.74a	0.68a	1.50	2.39b	0.85	0.51
SI	2.53b	36.56b	0.62b	1.42	2.34b	0.92	0.51
Sex							
Male	2.65	37.78	0.66	1.46	2.35	0.94a	0.48a
Female	2.65	37.40	0.66	1.46	2.45	0.85b	0.56b
Slaughter weight (kg)							
30	2.10a	35.68a	0.52a	1.17a	1.78a	0.65a	0.33a
35	2.47b	36.85a	0.61b	1.37b	2.19b	0.80b	0.44a
40	2.84c	37.68b	0.72c	1.56c	2.67c	1.03c	0.58a
45	3.09d	39.53b	0.76c	1.68c	2.85c	1.06c	0.67a
Regression	30.459+0.005*LW ²	30.7316+0.0052*LW ²	0.247+0.012*LW	0.299+0.336*LW	0.318+0.062*LW	0.232+0.019*LW	-0.143+0.020*LW

Means with different letters in the same column are significantly different (P<0.05), using Duncan test.

T × SI - ½ Texel × ½ Santa Ines; I × SI - ½ Ile de France × ½ Sant Ines, SI - pure Santa Ines; LW - live weight.

There was a high negative correlation between birth weight and days on creep feeding, indicating that a heavier animal at birth spent less time to reach weaning weight (Table 8). Villas Bôas et al. (2003) found that creep feeding led to lambs being weaned earlier with improved killout and carcass traits. The mean negative correlation between

days to slaughter and weight gain on finishing indicates that slower growing animals spent more time to finishing.

Hot carcass weight had medium positive correlations with leg perimeter (0.48) and shoulder weight (0.32), and half-carcass weight had medium positive correlations with skin, fat score, head, shoulder, leg, back and belly weights

Table 6 - Mean comparison for non carcass cuts in Santa Ines and crossbred lambs slaughtered at different weights

Source of variation	Thoracic organs (kg)	Abdominal organs (kg)	Rumen (kg)	Intestines (kg)	Head (kg)	Skin (kg)
Genetic group						
Texel × Santa Ines	0.92	1.15	3.56a	3.91a	2.24a	3.24a
Ile de France × Santa Ines	0.91	1.16	4.55b	2.75b	1.82b	3.32a
Santa Ines	0.91	1.27	4.20b	2.79b	1.77b	2.49b
Sex						
Male	0.94	1.07a	4.70a	2.86	1.95a	3.11a
Female	0.88	1.36b	3.77b	3.02	1.77b	2.75b
Slaughter weight (kg)						
30	0.82a	0.93a	4.13	2.43a	1.54a	2.30a
35	0.88b	1.13b	4.18	2.83b	1.77b	2.75a
40	0.98b	1.34b	4.20	2.92b	1.83b	2.93b
45	0.96b	1.40b	4.40	3.41b	2.20c	3.54b
Regression	$-2.112+0.165*LW-0.002*LW^2$	$0.565+0.009*LW$	NS	$0.759+0.064*LW$	$0.167+0.049*LW$	$0.812+0.002*LW^2$

Means with different letters in the same column are significantly different ($P<0.05$), using Duncan test. NS - not significant.

Table 7 - Means for measurements taken on the *longissimus dorsi* muscle in purebred and crossbred sheep slaughtered at different weights

Source of variation	Rib Eye S (cm ²)	Depth (cm)	Height (cm)	Fat (mm)
Genetic group				
Texel × Santa Ines	14.87a	5.93	3.62a	3.50a
Ile de France × Santa Ines	13.08b	5.69	3.06b	3.40ab
Santa Ines	12.26b	5.67	3.06b	3.25b
Sex				
Male	12.87	5.81a	3.08a	2.51a
Female	13.19	5.61b	3.21b	4.32b
Slaughter weight (kg)				
30	11.36a	5.45a	2.86a	2.90a
35	11.93a	5.60a	3.13b	3.06b
40	13.82b	5.77b	3.18b	3.54b
45	14.65b	5.97b	3.33b	3.96b
Regression	$2.28+0.30*LW$	$2.5349+0.0008*LW^2$	$3.734+0.053*LW$	$0.3170+0.0001*LW$

Means with different letters in the same column are significantly different ($P<0.05$), using Duncan test.

Rib Eye S - rib eye area using squared transparent paper; Depth - distance between lateral extremities of the *longissimus dorsi* at the 12th rib; Height - distance between inferior and superior extremities of rib eye at the 12th rib; Fat - fat thickness measured using a paquimeter.

Table 8 - Correlations between growth traits in lambs

	BW	WW	Days WW	WGC	WG All	SW	Days SW
WW	-0.75						
Days	-0.85	0.80					
WGC	-0.27	-0.56	-0.23				
WG All	0.86	-0.79	-0.72	-0.02			
SW	-0.85	0.53	0.70	0.29	-0.82		
Days SW	-0.60	0.65	0.80	-0.26	-0.52	0.50	
WG SW	0.72	-0.97	-0.75	0.48	0.82	-0.55	-0.57

BW - birth weight; WW - weaning weight; Days WW - days to weaning; Days SW - days to slaughter; WG All - weight gain from birth to slaughter; WGC - weight gain in creep feeding; SW - slaughter weight.

Table 9 - Correlations between qualitative and quantitative carcass traits in lambs

	HCW	CCW	CWH	Skin	TO	AO	Fat	Rum	Intestines	Head	Leg	Legcir	Back	Shoulder	Back	Neck
CCW	-0.24															
CWH	0.20	-0.50														
Skin	-0.18	-0.46	0.31													
TO	-0.35	-0.36	-0.01	0.75												
AO	-0.29	-0.17	-0.19	0.30	0.84											
Fat	-0.60	0.26	0.43	0.01	-0.15	-0.24										
Rum	-0.30	0.10	-0.55	0.52	0.60	0.41	-0.31									
Intestines	-0.28	-0.23	-0.13	0.10	-0.01	0.53	-0.40	0.08								
Head	0.07	-0.67	0.56	0.59	0.61	0.47	-0.05	0.02	0.49							
Leg	0.28	-0.15	0.02	0.10	0.21	0.25	-0.32	0.19	0.23	0.26						
Legcir	0.48	0.54	-0.46	-0.82	-0.76	-0.44	-0.34	-0.25	-0.35	-0.68	-0.02					
Back	-0.63	0.18	-0.47	0.13	-0.04	-0.17	0.99	-0.26	0.10	0.05	-0.31	-0.45				
Shoulder	0.33	-0.84	0.55	0.74	0.40	0.02	-0.26	0.08	-0.15	0.67	0.14	-0.58	-0.16			
Ribs	-0.52	0.30	-0.65	0.38	0.68	0.63	-0.16	0.85	0.24	-0.02	0.11	-0.26	-0.12	-0.20		
Neck	-0.08	-0.29	0.25	-0.18	0.33	0.67	0.03	-0.33	0.91	0.44	0.18	-0.22	0.04	-0.08	-0.08	
Belly	0.09	-0.65	0.51	0.93	0.63	0.19	-0.07	0.28	-0.02	0.69	0.15	-0.76	0.04	0.89	0.09	-0.10

CCW - cold carcass weight; HCW - hot carcass weight; CWH - half carcass weight; Skin - skin weight; Legcir - leg circumference; Fat - fat score; TO - thoracic organs; AO - abdominal organs; Rum - rumen weight.

Table 10 - Correlations between morphological and eye muscle measures in lambs

	FAT	EMA	MW	MH	SH	TC	CC	BS	EMAUS
EMA	-0.30								
MW	-0.33	0.27							
MH	0.92	-0.48	-0.21						
SH	-0.68	-0.51	-0.03	-0.49					
TC	-0.02	-0.90	-0.35	0.13	0.69				
CC	0.53	0.03	-0.19	0.52	-0.52	-0.40			
BS	0.81	0.32	-0.07	0.66	-0.97	-0.57	0.61		
EMAUS	-0.48	0.31	0.98	-0.37	0.03	-0.30	-0.53	-0.17	
FTUS	-0.71	0.32	0.00	-0.78	0.45	0.12	-0.85	-0.59	0.85

FAT - fat score; EMA - eye muscle area; MW - muscle width; MH - muscle height; SH - shoulder height; TC - thoracic circumference; CC - cannon bone circumference; BS - body score; EMAUS - eye muscle area with ultrasound; FTUS - fat thickness with ultrasound.

($r = 0.31$; $r = 0.43$; $r = 0.56$; $r = 0.55$; $r = 0.02$; $r = 0.47$ and $r = 0.51$), respectively (Table 9). Thoracic organ weight had a positive correlation with abdominal organ weights, rumen, leg, head, rib and belly weight, while abdominal organ weight had high positive correlation with rib and neck weights.

Fat thickness (Table 10) and eye muscle area decreased with an increase in morphometric measures, indicating that larger animals are later-developing in terms of carcass finishing than smaller animals. For Prado et al. (2004), the eye muscle area is used as an indicator of muscle development.

Conclusions

The Texel and Ile de France crossbred lambs are superior to the pure Santa Ines in terms of carcass quality. The ideal slaughter weight is between 35 and 40 kg for Santa Ines, while females deposit fat earlier than males, meaning their slaughter weight should be lower.

Acknowledgments

To CNPq (INCT-IGS), Finatec and FAPDF for financial support.

References

- ARNOLD, A.M.; MEYER, H.H. Effects of gender, time of castration, genotype and feeding regimen on lamb growth and carcass fatness. *Journal of Animal Science*, v.66, p.2468, 1988.
- BRASAL, T.L.; BOCCARD, R. Efectos de los tratamientos antemortem sobre la calidad de la canal y la carne de cordero. INIA, *Série Producción Animal*, v.8, p.97-125, 1977.
- CARDELLINO, R.A. Sistemas de produção de carne ovina utilizando cruzamentos. In: SIMPÓSIO PAULISTA DE OVINO-CULTURA, 1989, Botucatu. *Anais...* Botucatu, 1989. p.97.
- COSTA, R.G.; MEDEIROS, A.N.; MADRUGA, M.S. et al. Rendimento de vísceras para "buchada" em caprinos Saanen alimentados com diferentes níveis de volumoso e concentrado. In: SIMPÓSIO INTERNACIONAL DE CAPRINOS E OVINOS DE CORTE, 2003, João Pessoa. *Anais...* João Pessoa: Emepa, 2003. p.663-666.
- FURUSHO-GARCIA, I.F.; PEREZ, J.R.O.; BONAGURIO, S. et al. Estudo dos cortes de carcaça de cordeiros da raça Santa Ines puros e cruzas Santa Ines com Texel, Ile de France e Bergamácia. *Revista Brasileira de Zootecnia*, v.33, p.453-462, 2004.
- HUIDOBRO, F.R.; VILLAPADIerna, A. *Estudios sobre crecimiento y desarrollo en corderos de raza Manchega*. 1992. 191p. Thesis (Doctoral) - Universidad Complutense, Madrid.
- JUÁREZ, M.; HORCADA, A.; ALCALDE, M.J. et al. Meat and fat quality of un weaned lambs as affected by slaughter weight and breed. *Meat Science*, v.83, p.308-313, 2009.
- LAMBE, N.R.; NAVAJAS, E.A.; SCHOFIELD, C.P. et al. The use of various live animal measurements to predict carcass and meat quality in two divergent lamb breeds. *Meat Science*, v.80, p.1138-1149, 2008.
- LANDIM, A.V.; MARIANTE, A.S.; McMANUS, C. et al. Características quantitativas da carcaça, medidas morfométricas e suas correlações em diferentes genótipos de ovinos. *Ciência Animal Brasileira*, v.8, p.665-676, 2007.
- LANDIM, A.V.; CASTANHEIRA, M.; FIORAVANTI, M.C.S. et al. Physical, chemical and sensorial parameters for lambs of different groups, slaughtered at different weights. *Tropical Animal Health and Production*, v.43, p.1089-1096, 2011a.
- LANDIM, A.V.; CARDOSO, M.T.M.; CASTANHEIRA, M. et al. Fatty acid profile of hair lambs and their crossbreds slaughtered at different weights. *Tropical Animal Health and Production*, v.43, p.1561-1566, 2011b.
- MACEDO, A.F.M.; SIQUEIRA, E.R.; MARTINS, N.E. Análise econômica da produção de carne de cordeiros sob dois sistemas de terminação: pastagem e confinamento. *Ciência Rural*, v.30, p.677-680, 2000.
- MARTINS, A.R.V. *Utilização de dejetos suínos em dietas de ovinos em sistema de confinamento*. 1997. 51f. Dissertação (Mestrado em Zootecnia) - Universidade Federal de Lavras, Lavras.
- McMANUS, C.; MIRANDA, R.M. Comparação de ovinos Santa Inês e Bergamácia no Distrito Federal. *Revista Brasileira de Zootecnia*, v.26, p.1055-1059, 1997.
- MENDONÇA, G.; OSÓRIO, J.C.; OLIVEIRA, N.M. et al. Morfologia, características da carcaça e componentes do peso vivo em borregos Corriedale e Ideal. *Ciência Rural*, v.33, p.351-355, 2003.
- MORENO, G.M.B.; SILVA SOBRINHO, A.G.; ROSSI, R.C. et al. Desempenho e rendimentos de carcaça de cordeiros Ile de France desmamados com diferentes idades. *Revista Brasileira de Saúde e Produção Animal*, v.11, p.1105-1116, 2010.
- OSÓRIO, J.C.S.; OSÓRIO, M.T.M.; JARDIM, P.O.C. *Métodos para avaliação de carne ovina "in vivo" na carcaça e na carne*. Pelotas: Ed. UFPEL. 1998. 107p.
- OSÓRIO, J.C.S.; OSÓRIO, M.T.M. Sistemas de avaliação de carcaça no Brasil. In: SIMPÓSIO MINEIRO DE OVINO-CULTURA: PRODUÇÃO DE CARNE NO CONTEXTO ATUAL, 1., 2001, Lavras. *Anais...* Lavras: Universidade Federal de Lavras, 2001. p.198.
- OSÓRIO, J.C.S.; OLIVEIRA, N.M.; OSÓRIO, M.T.M. et al. Produção de carne em cordeiros Border Leicester com ovelhas Corriedale e Ideal. *Revista Brasileira de Zootecnia*, v.31, p.1469-1480, 2002.
- PAIM, T.P.; CARDOSO, M.T.M.; BORGES, B.O. et al. Estudo econômico da produção de cordeiros cruzados confinados abatidos em diferentes pesos. *Ciência Animal Brasileira*, v.12, p.48-57, 2011.
- PÉREZ, P.; MAINO, M.; MORALES, M.S. et al. Gender and slaughter weight effects on carcass quality traits of suckling lambs from four different genotypes. *Small Ruminant Research*, v.70, p.124-130, 2007.
- PILAR, R.C.; PÉREZ, J.R.O.; SANTOS, C.L. Manejo reprodutivo da ovelha recomendações para uma parição a cada 8 meses. *Boletim Agropecuario*, v.50, p.1-28, 2002.
- PINHEIRO, R.S.B. *Aspectos quantitativos da carcaça e qualitativos da carne de ovinos de diferentes categorias*. 2006. 115f. Dissertação (Mestrado em Zootecnia) - Faculdade de Ciências Agrárias e Veterinárias/Universidade Estadual Paulista, Jaboticabal.
- PINHEIRO, R.S.B.; SILVA SOBRINHO, A.G.; SOUZA, H.B.A. et al. Informações nutricionais de carnes ovinas em rótulos comerciais, comparativamente às obtidas em análises laboratoriais. *Ciência e Tecnologia de Alimentos*, v.27, p.376-381, 2007.
- PIRES, C.C.; CARNEIRO, R.M.; MULLER, L. et al. Avaliação da carcaça e componentes do peso vivo, de cordeiros de parto simples desmamados, parto simples não desmamados e de parto duplo desmamado. *Revista Brasileira de Zootecnia*, v.12, p.93-97, 2006.
- PIRES, C.C.; MÜLLER, L.; TONETTO, C.J. et al. Influência do tipo de parto e do sexo no desempenho e nas características de carcaça de cordeiros cruza Ile de France x Texel. *Revista Ceres*, v.58, p.432-437, 2011.
- PRADO, C.S.; PÁDUA, J.T.; CORREA, M.P.C. et al. Comparação de diferentes métodos de avaliação da área de olho de lombo e cobertura de gordura em bovinos de corte. *Ciência Animal Brasileira*, v.5, p.141-149, 2004.
- RIBEIRO, L.A.O.; GREGORY, R.M.; MATTOS, R.C. Prenhez em rebanhos ovinos do Rio Grande do Sul. *Ciência Rural*, v.32, p.637-641, 2002.
- ROQUE, A.P.; OSÓRIO, J.C.S.; JARDIM, P.O. et al. Produção de carne em ovinos de cinco genótipos. Desenvolvimento relativo. *Ciência Rural*, v.29, p.549-553, 1999.
- ROSA, G.T.; PIRES, C.C.; SILVA, J.H. et al. Proporções e coeficientes de crescimento dos não componentes da carcaça de cordeiros e cordeiras em diferentes métodos de alimentação. *Revista Brasileira de Zootecnia*, v.31, p.2290-2298, 2002.
- SANTANA, A.F. Correlação entre peso e medidas corporais em ovinos Jovens da Raça Santa Ines. *Revista Brasileira de Saúde e Produção Animal*, v.1, p.74-77, 2001.
- SILVA SOBRINHO, A.G. Aspectos quantitativos e qualitativos da produção de carne ovina. In: REUNIÃO ANUAL DA SOCIEDADE BRASILEIRA DE ZOOTECNIA, 38., 2001,

- Piracicaba. **Anais...** Piracicaba: Sociedade Brasileira de Zootecnia, 2001. p.425-446.
- SILVA, L.F.; PIRES, C.C.; ZEPPENFELD, C.C. et al. Crescimento de regiões da carcaça de cordeiros abatidos com diferentes pesos. **Ciência Rural**, v.30, p.481-484, 2000.
- TAYLOR, C.S. Use of genetic size scaling in evaluation of growth. **Journal of Animal Science**, v.61, p.1103-1107, 1985.
- VILLAS BÔAS, A.S.; ARRIGONI, M.D.B.; SILVEIRA, A.C. et al. Idade à desmama e manejo alimentar na produção de cordeiros superprecoces. **Revista Brasileira de Zootecnia**, v.32, p.1969-1980, 2003.
- ZUNDT, M.; MACEDO, F.A.F.; MARTINS, E.N. et al. Desempenho de cordeiros alimentados com diferentes níveis de proteína. **Revista Brasileira de Zootecnia**, v.31, p.1307-1314, 2001.